arXiv:1208.2660v2 [hep-th] 24 Sep 2012

A Numerical Simulation of Chern-Simons Inflation

Annie Preston[1], **David Garrison**[2] **and Stephon Alexander**[3]

[1] Department of Physics, Harverford College, Harverford, PA 19041
[2] Physics Department, University of Houston Clear Lake, Houston, TX 77058
[3] Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755

Abstract. In this work, we present results of numerical simulations of the Chern-Simons Inflation Model proposed by Alexander, Marciano and Spergel. According to this model, inflation begins with a fermion condensate interacting with a gauge field. Crucial to the success of this mechanism is the assumption that the Chern-Simons interaction would drive energy from the initial random spectrum into a narrow band of frequencies at superhorizon scales. In this work we numerically confirm this expectation. These gauge fields, when combined with the Friedmann equations, were broken into a system of hyperbolic equations and numerically simulated with a novel relativistic MHD code. We show that the amplification of horizon sized gauge fields produces the conditions to cause cosmological inflation and that the onset of inflation are robust against certain fine tunings in the initial conditions.

1. Introduction and Motivation

The paradigm of the early universe is based on the phenomenon of cosmic inflation (Guth 1981). It is well known by now that inflation resolves most of the problems of the standard big-bang scenario, such as the horizon problem. Moreover, inflation predicts a nearly scale-invariant power spectrum of density perturbations which is observed in the CMB. Although, there many successes of inflation, there are hundreds of scalar field models of inflation which have proven difficult to distinguish with data. Moreover, scalar field driven inflation are frought with conceptual and technical issues. Recently Chen and Wang demonstrated that the primordial power-spectrum in scalar field inflation have anomalous sensitivity to high energy physics (Chen & Wang 2012). These issues have motivated the authors (Alexander et al. 2011) to provide an inflationary mechanism that is driven by an interaction between vector and spinor fields, otherwise known as Chern-Simons inflation.

Alexander, Marciano, and Spergel propose a model in which the early universe is dominated by a gauge field that interacts with a fermion current. This interaction results in an equation of state with consistently negative pressure, the condition needed for inflation (Alexander et al. 2011). In this model, the gauge field begins as a random, white noise spectrum; the authors assume that this evolves into a spectrum of superhorizon modes. Because of the complexity of the differential equations, they were not able to show this evolution analytically. The purpose of this investigation is to determine numerically whether the gauge field transforms from a random spectrum into a spectrum of low-frequency modes during inflation.

Another goal of this project is to establish the robustness of the model. One criticism of inflation is that specific theories can focus too much on formalism and lack a clear connection to physical processes that actually could have occurred (Kinney 2009). Therefore, we aim to strengthen the physical interpretation of an inflationary theory while assessing its accuracy; we use a modified magnetohydrodynamic (MHD) code based on the Cactus framework to simulate the early universe. Though this is perhaps the first cosmological MHD inflation simulation, there have been many astrophysical MHD simulations, and there has been a wide range of other types of inflation simulations. For example, in (Dubois & Teyssier 2008), the authors use an MHD solver to simulate the environment within galaxy clusters. Examples of other inflationary models that have been simulated include a back-reacting cosmology with inhomogeneous initial conditions (Kurki-Suonio et al. 1987), pre-Big Bang inflation (Maharana et al. 1998), and an examination of topological defects due to inflation (Kasuya & Kawasaki 1998). Our work presents another method of testing new inflationary theories in an accessible way, and introduces the concept of treating the early universe as an MHD like system.

In Section 2, we provide necessary mathematical background, then we explain motivations for all possible mechanisms of inflation. In Section 3, we describe the analytical and computational methods used to simulate the specific inflationary theory from (Alexander et al. 2011). In Section 4, we present the results of the simulation, and in Section 5, we discuss the results and future research directions.

2. Inflation Models

The accelerated expansion during inflation requires that the universe be dominated by a negative-pressure component. In many inflationary models, a scalar field dominates the early universe and provides the necessary negative-pressure equation of state. The following discussion is closely based on (Ryden 2003). This scalar field, $\phi(\vec{r}$, t), is known as the inflaton field. It has a potential V(ϕ) that depends on the temperature of the radiation dominating the universe prior to inflation (Montani et al. 2011). The energy density and pressure of the inflaton field are

$$\epsilon_\phi = \frac{1}{2}\frac{1}{\hbar c^3}\dot{\phi}^2 + V(\phi) \tag{1}$$

$$P_\phi = \frac{1}{2}\frac{1}{\hbar c^3}\dot{\phi}^2 - V(\phi) \tag{2}$$

If the change in the inflaton field over time is very slow:

$$\dot{\phi}^2 \ll \hbar c^3 V(\phi) \tag{3}$$

then we have

$$\epsilon_\phi \approx -P_\phi \tag{4}$$

which meets the requirement for negative-pressure domination. We can see here that if there is a period in which $\dot{\phi}$ is small, and $V(\phi)$ is large enough to dominate the energy density of the universe, then the inflaton field can cause exponential expansion. This model is known as slow roll inflation. However, it is not immediately clear what these conditions physically correspond to in the early universe. We can determine the necessarily small value of $\dot{\phi}$ by considering the inflaton field′s fluid equation:

$$\dot{\epsilon_\phi} + 3H(t)(\epsilon_\phi + P_\phi) = 0 \tag{5}$$

This gives

$$\ddot{\epsilon_\phi} + 3H(t)\dot{\epsilon_\phi} = -\hbar c^3 \frac{dV}{d\phi}, \tag{6}$$

which is an equation of motion in which the accelerating force is -$\frac{dV}{d\phi}$ and the “frictional” force, proportional to speed, is 3H(t). This force, known as the “Hubble friction,” slows the change in the inflaton field until its potential, $V(\phi)$, is at a minimum. When the acceleration $\ddot{\phi}$ reaches zero, we have

$$\dot{\phi} = -\frac{\hbar c^3}{3H}\frac{dV}{d\phi}. \tag{7}$$

We know that the requirement in Equation 3 is necessary for the inflaton field to cause inflation, so imposing that requirement gives

$$(\frac{dV}{d\phi})^2 \ll \frac{9H^2V}{\hbar c^3}. \tag{8}$$

When the potential energy of the inflaton dominates, which is necessary for inflation, the Hubble parameter is

$$H = (\frac{8\pi G\epsilon_\phi}{3c^2})^{1/2} = (\frac{8\pi GV}{3c^2})^{1/2} \tag{9}$$

and our requirement becomes

$$(\frac{E_P}{V}\frac{dV}{d\phi})^2 \ll 1, \tag{10}$$

where E_P is the Planck energy. Note that during inflation, the Hubble parameter is constant. We have found that the inflaton′s potential must satisfy the equation above, and that its energy density must be large enough to dominate that of the universe, in order for exponential expansion to occur. These conditions imply a nearly constant energy density of the inflaton field: $\rho \sim \dot{\phi}^2 + V(\phi)$; this requires its kinetic energy to be insignificant compared to its potential energy.

To see how inflation solves the Flatness problem, consider yet another version of the Friedmann equation:

$$(\Omega^{-1} - 1)\rho a^2 = -\frac{3\kappa c^2}{8\pi G} \tag{11}$$

If the scale factor, a, increases exponentially while the energy density, ρ, stays nearly constant, Ω must approach 1, since the right side of the equation is constant. During inflation, the energy density of the universe is dominated by that of the inflaton field, which remains constant just as the energy density of dark energy does. Therefore, inflation drives the universe towards flatness. Because the Hubble parameter is fixed during inflation, the curvatures immediately before and after inflation are related by

$$\Omega_f - 1 = (\Omega_i - 1)e^{-2N} \tag{12}$$

where N is the number of e-folds. According to the Standard Cosmological Model, the quantity Ω - 1 has increased by a factor of about 10^{60} since the end of inflation; therefore, the current observational constraint on $|\Omega_0 - 1|$ gives

$$|\Omega_0 - 1| = 10^{60}|\Omega_i - 1|e^{-2N} \lesssim 10^{-2}. \tag{13}$$

The minimum number of e-folds needed to resolve the flatness paradox is then

$$N \gtrsim 71 + \frac{1}{2}|\Omega_i - 1|. \tag{14}$$

We see that even with significant initial deviations from flatness, enough e-folds of inflation efficiently flatten the universe.

2.1. Chern-Simons Inflation

Because these scalar field models are widely found and difficult to distinguish from one another, Alexander, Marciano and Spergel suggest an alternate model in which a gauge field interacts with fermions in the early universe to produce an effective scalar field that generates inflation (Alexander et al. 2011).

In this gauge field model, we have an energy density

$$\rho = \frac{E^2 + B^2}{2a^4} + |A \cdot \mathcal{J}| \tag{15}$$

where the gauge field has an electric field, $\vec{E} \equiv \dot{\vec{A}}$, and a magnetic field, $\vec{B} \equiv \vec{\nabla} \times \vec{A}$. We can see that this energy density has an electromagnetic component, which scales as a^{-4} , as well as a gauge field-fermion interaction. As shown in (Alexander et al. 2011), if the second term dominates over the first term, while remaining nearly constant, it is possible for inflation to occur. This will give us an energy density $\rho = |A \cdot \mathcal{J}|$ and pressure P = -$|A \cdot \mathcal{J}|$, corresponding to an equation of state w = -1, which is sufficient to cause inflation.

Just as we gained insight through the scalar field′s equation of motion for the general case, we will consider the equation of motion of the gauge field in this case. Its action is:

$$\begin{aligned} S = \int_{M_4} d^4x\sqrt{-g}(\frac{M_P^2 R}{8\pi} - \frac{1}{2}\partial_\mu\theta\partial^\mu\theta + \xi R\theta^2 - \frac{1}{4}Tr[F_{\alpha\beta}F^{\alpha\beta}] \\ + \frac{\theta}{4M_*}Tr[F_{\alpha\beta}F^{\alpha\beta}] + qTr[A_\mu \mathcal{J}_5^\mu]) \end{aligned} \tag{16}$$

By varying this action with respect to the gauge field, we find the gauge field′s equation of motion. In terms of Fourier modes, this equation of motion is:

$$\ddot{A}_h + k^2 A_h = -hkA_h\frac{\dot{\theta}}{M_*} + a^4\mathcal{J}_h \tag{17}$$

where the fermion currents are

$$\sqrt{2}\mathcal{J}_h = \mathcal{J}_1 + hi\mathcal{J}_2 = (\bar{\psi}(\gamma_1 + hi\gamma_2)\gamma_5\psi)/a. \tag{18}$$

Alexander et al. show that the fermion currents have a background solution in which they scale as J_0/a . Since they are being suppressed by a factor of a, we can define a set of constants J_h where $J_h = \mathcal{J}_h/a$ (Alexander et al. 2011). The equation of motion for the gauge field is now

$$\ddot{A}_h + k^2 A_h = -hkA_h\frac{\dot{\theta}}{M_*} + a^3 J_h \tag{19}$$

According to the solutions for the gauge field (Alexander et al. 2011), a value of $k < \frac{\dot{\theta}}{M_*}$, corresponding to long-wavelength modes, will result in its exponential growth. However, the presence of short wavelength modes, $k > \frac{\dot{\theta}}{M_*}$, will result in continued linear growth due to back-reaction of the gravitational field.

Because the universe was a plasma before last scattering, we look to plasma physics for an analogy. Specifically, we consider magnetohydrodynamics (MHD), the study of electrically conducting fluids. When fluids in motion are electrically conducting, currents in the fluid produce magnetic fields that induce forces and thus change the dynamics of the system when the system becomes turbulent (Shebalin 2010). This kind of turbulence is analogous to the dynamical system changes that could be capable of generating inflationary behavior. Computer codes that simulate MHD use the equations of fluid dynamics, along with Maxwell′s equations, to analyze the dynamics of a plasma system. Simulations such as these are commonly known to lead to what is referred to as an "inverse energy cascade". In a hydrodynamic turbulent system, as a system evolves in time, energy tends to "cascade" from the longest wavelength modes to the shortest wavelength modes as turbulent eddies get smaller and smaller. In MHD systems, the reverse happens. Energy cascades inversely from high frequency modes to low frequency modes.

Modifying an MHD code by adding the coupled partial differential equations that describe the very early universe could allow us to see the resulting exponential increase and then "calming down" of the scale factor. Additionally, a simulation of this type will allow us to see directly the physical effects of fine-tuning the initial conditions for these coupled differential equations. We can input the equations of motion for the fields that are interacting to generate inflation, along with the Friedmann equations, which describe the expansion of space. The Friedmann equations will allow us to relate the evolving pressure and density of the system to the evolution of the scale factor.

3. Methods and Equations

For the numerical calculation, we use natural units but later evaluate the data in terms of SI units so that the results can be easily compared to the established values. The relevant equation from (Alexander et al. 2011) is the equation of motion for the gauge field, A:

$$\ddot{\vec{A}} - \nabla^2 \vec{A} + \frac{\dot{\theta}}{M_*}(\vec{\nabla} \times \vec{A}) - a^4 \mathcal{J}_i = 0 \tag{20}$$

Here, $\mathcal{J}$ is the fermion current and M_* is a scale mass associated with the theory (Alexander et al. 2011). In comoving coordinates this equation becomes:

$$a(t)\frac{d^2\vec{A}}{dt^2} + \frac{da}{dt}\frac{d\vec{A}}{dt} = \vec{J}a(t)^3 + \frac{1}{a(t)}\nabla^2\vec{A} - a(t)^2\frac{\dot{\theta}}{M_*}\vec{\nabla} \times \vec{A} \tag{21}$$

In order to use this in our code, we separated the equations of motion for the gauge field and the Friedman equations into a system of first order in time differential equations.

$$\frac{d\vec{A}}{dt} = \frac{\vec{Z}}{a} \tag{22}$$

$$\frac{d\vec{Z}}{dt} = \vec{J}a^3 + \nabla^2\vec{A}/a - a^2\frac{\dot{\theta}}{M_*}(\vec{\nabla} \times \vec{A}) \tag{23}$$

$$\frac{d\vec{a}}{dt} = aH \tag{24}$$

$$\frac{d\vec{H}}{dt} = \frac{8\pi}{3}\bar{\rho}/a - H^2 \tag{25}$$

The average energy density and pressure are calculated based on equation 15 as $\bar{\rho} = \frac{1}{N}\sum_{k=1}^{N} |A_k \cdot \mathcal{J}_k|$ and $\bar{P} = -\frac{1}{N}\sum_{k=1}^{N} |A_k \cdot \mathcal{J}_k|$. Here N represents the total number of grid points in the computational domain. The scale factor and Hubble parameter therefore depend on the average energy density and not the local field dynamics. Because of this, the gravitational back-reaction of the Chern-Simons gauge field on the field equations is assumed negligible.

Once the equations were established we input the initial conditions assuming that the current was constant in time (it should be slowly varying) and that the gauge field was composed of a random (white noise) spectrum. In order to generate the initial gauge field, we used a random number generator to create a random spectrum with amplitude up to the calculated maximum amplitude, A_0, in each direction. The magnitude of the gauge field was then held equal to the initial amplitude A_0. The initial value for the other variables is given below.

$$|\vec{A}_0| = 10^{-5} M_P \tag{26}$$

$$\vec{J} = 10^{-20} M_P^3 (\hat{x} + \hat{y} + \hat{z}) \tag{27}$$

$$\frac{\dot{\theta}}{M_*} = 2.18 \times 10^{-5} M_P \tag{28}$$

$$a_0 = 1.0 \tag{29}$$

$$H_0 = \sqrt{\frac{8\pi}{3} |\vec{A}_0 \cdot \vec{J}|} \tag{30}$$

$$\vec{Z}_0 = H_0 a_0 \times random\ number(-1, 1) \tag{31}$$

The code was then run on the University of Houston Texas Learning & Computation Center′s Xanadu cluster using a variety of time-steps, grid sizes and resolutions in order to obtain consistent results. We ran the code using resolutions ranging from 34^3 to 132^3 and grid sizes ranging from 10^4 to 10^7 cubed. These grid sizes correspond to the size of the horizon throughout the simulation. We adjusted the courant size from a low of 0.001 to a high of 0.5. These runs utilized anywhere from 16 to 48 processors and the runs lasted anywhere from 2 hours to 7 days. We also ran the code using three different differencing schemes including 2nd order finite differencing, 4th order finite differencing and Fourier spectral differencing.

Because the initial units were entered as Planck units, we assumed that the physical grid (horizon) size corresponded to Planck length and the timing output could be interpreted as Planck time. The output was analyzed using several tools including ygraph and Pro Fit (www.quansoft.com). We used both Pro Fit and a custom routine, which utilized FFTW, to perform a spectral analysis of the gauge field.

4. Results

Assuming that the inflationary period begins when the second derivative of the scale factor is positive, we find that inflation occurs in our simulation whenever the grid size is on the order of 10^6 or larger cubed. In these cases, we find that inflation consistently begins around 9.11×10^{-38} s. This roughly agrees with the established inflation theory. The scale factor continues to increase exponentially until the end of the evolution.



Figure 1. The scale factor before and during the inflationary period of the simulation.

The power spectrum of the gauge field clearly shows the development of an inverse energy cascade during the inflationary phase. The initial power spectrum showed a flat (random noise) energy distribution with energy variations of up to 2 orders of magnitude. As the system evolved, the lowest order modes became more energetic relative to all other parts of the spectrum with over 5 orders of magnitude more energy than the highest frequency modes. Later in the evolution, the spectrum appears to have flattened out as energy returned to the high-frequency modes and variations in the power spectrum became smaller. Here high frequency modes correspond to small scale inhomogeneities so during inflation, the system appears to become more homogenous in local regions of space-time.

For computational grids with a horizon size of 10^6 planck lengths cubed or larger, we obtained results consistent with inflation, regardless of the step-size, differencing method or other numerical factors. For smaller grids, however, this exponential expansion does not take place. This implies that the inflation process requires wavelengths on the order of 10^6 planck lengths. This is most likely a function of our initial conditions for the gauge field, current and Chern-Simons interaction terms.



Figure 2. The Power Spectral Density of the Gauge Field at different planck times. The vertical axis shows the log of the deviation from the mean. The horizontal axis shows frequency in planck units.

5. Discussion and Conclusions

The simulation presented here clearly supports much of the hypotheses proposed by Alexander, Marciano and Spergel: the evolution equations given drive the energy into a narrow band of modes and cause the universe to expand exponentially. Though the theory does not predict an initial start time for inflation, our value of about $t_i = 10^{-37}$ s is consistent with the time predicted by the most general models of inflation (Weinberg 2008). This suggests some universality in the initial conditions required for inflation, regardless of the model; in the future, we can study this question analytically for this model. However, we do not see inflation end in our simulation, so we cannot verify the validity of the mechanism for ending inflation, or predict a stopping time numerically.

The power spectrum in Figure 2 supports the hypothesis that an initially random gauge field spectrum becomes concentrated in low frequencies during inflation. This analysis allows us to see the progression from an almost completely random spectrum near the beginning to the energy being concentrated in a narrow band of low-frequency modes during the inflation event. In addition to the dominance of low-frequency modes, the higher modes are suppressed during inflation; the beating pattern probably results from imperfections in the random number generator. This reflects the behavior predicted in (Alexander et al. 2011). Though we were not able to see the end of inflation in this simulation, the latest time power spectrum suggests that inflation is in fact ending. The dominance of low-frequency modes has subsided, and we are mostly left with random noise behind a flat power spectrum, indicating that the end of inflation is near.

It is interesting that exponential growth does not occur for horizon sizes that are too small. This could be due to the unavailability of very long-wavelength (low frequency) modes for the gauge field. In the future, accurate measurements of these frequencies could allow us to verify that they are superhorizon modes. However, the most logical next step in this research will involve improving the dynamics of the gauge field and fermion current to better predict the length and end of inflation. This may involve changes to the evolution equations for the gauge field or fermion current during the inflation event which may also have an impact on the field dynamics at the sub-horizon level.

Acknowledgments

Thanks to the Louis Green Fund at Haverford College for funding travel to Houston, and to the University of Houston's Texas Learning and Computing Center for access to and use of the Xanadu computing cluster. Thanks to Peter Love for time spent on edits and helpful suggestions. SA was supported by the NSF CAREER grant and he thanks the Princeton physics department for their hospitality.

References

Alexander S, Marciano A & Spergel D 2011 *ArXiv e-prints* **hep-th**(1107.0318).
Chen X & Wang Y 2012 *ArXiv e-prints* **hep-th**(1207.0100).
Dubois Y & Teyssier R 2008 *AAP* **482**, L13–L16.
Guth A H 1981 *Phys. Rev. D* **23**(347).
Kasuya S & Kawasaki M 1998 *Phys. Rev. D* **58**, 083516.
URL: *http://link.aps.org/doi/10.1103/PhysRevD.58.083516*
Kinney W H 2009 *ArXiv e-prints* **astro-ph**(0902.1529).
Kurki-Suonio H, Centrella J, Matzner R A & Wilson J R 1987 *Phys. Rev. D* **35**, 435–448.
URL: *http://link.aps.org/doi/10.1103/PhysRevD.35.435*
Maharana J, Veneziano G & Onofri E 1998 *JHEP* **1998**(04), 004.
URL: *http://stacks.iop.org/1126-6708/1998/i=04/a=004*
Montani G, Battisti M V, Benini R & Imponente G 2011 *Primordial Cosmology* 1st, edn World Scientific New Jersey.
Ryden B 2003 *Introduction to Cosmology* 1st, edn Addison Wesley San Francisco.
Shebalin J 2010 *Physics of Plasmas* **17**.
Weinberg S 2008 *Cosmology* 1st, edn Oxford University Press Oxford.